

Redmond Linden

Tilia americana (Redmond)



Flower



Bark

General Attributes

Type	Deciduous Tree
Height	40 - 60 Feet
Spread	20 - 30 Feet
Form	Pyramidal
Utility Lines	Incompatible
Growth Rate	Medium
Life Expectancy	Medium
USDA Zone	4 - 7
Root Pattern	Deep Lateral

Flowers/Foliage/Fruits

Flower Color	Yellow
Flower Season	Summer
Fruit	Nut
Fruit Color	Tan
Fruit Season	Fall
Summer Texture	Medium
Winter Texture	Medium
Spring Foliage	Green
Summer Foliage	Green
Fall Foliage	Yellow
Winter Foliage	Not Applicable

Plant Community

Sun	Full Sun, Partial Sun
Orientation	North, South, East, West
Soil Texture	Sandy Loam to Clay Loam
Topography	Upland
Plant Community	Forest
Succession	Intermediate, Climax
Origin	North America

Soils

Salt Spray Tolerance	Sensitive
Soil Salt Tolerance	
Compaction Tolerance	Intolerant
Water Table	24 Inches
Drainage	Moderate
Flood Tolerance	
Drought Tolerance	
Moisture Regime	Moist, Wet
pH	6.0 through 8.0
Windbreak Group	

Redmond Linden

Tilia americana (Redmond)



Planting Ease

Spring Bareroot

Fall Bareroot

Spring Container

Easy

Fall Container

Easy

Spring Seed

Fall Seed

Maintenance

Formal

Low

Informal

Low

2,4-D Tolerance

Sensitive

Dicamba Tolerance

Sensitive

Picloram Tolerance

Sensitive

Clopyralid Tolerance

Sensitive

Artificial Light

Sulfur Dioxide

Tolerant

Ozone Tolerance

Tolerant

Hydrogen Flouride

Tolerant

Nitrogen Oxide

Moderate

Pests/Problems

Allergens

males 7, females 7

Invasiveness

Non-Invasive

Toxicity

Cold Injury

Moderately Susceptible

Storm Damage

Susceptible

Biological Control

Mechanical Control

Chemical Control

Miscellaneous

Wildlife Rating

Moderate

Insect Concerns

Minor

Disease Concern

Minor

Wildlife Concerns

Comments

Introduced by Plumfield Nurseries of Fremont, Nebraska in 1927. Originally listed as *T. x euchlora*. Reddish branches in winter and largest leaves of the lindens. Tests show slow to establish. Good street tree. Protect young bark from sunscald.



Dormant